



DISEASE PROTECTION FROM *PYTHIUM* AND *PHYTOPHTHORA*

- Labeled to be soil applied on many ornamental and turf species
- Effective control of oomycete (water mold) pathogens
- Can be applied through irrigation systems
- Compatible in tank mixes
- Curative + preventive activity
- No restricted entry interval under certain application methods

USE SITES - ORNAMENTALS

Christmas Tree Plantations	Greenhouses
Citrus (non-bearing) in nurseries & landscapes	Ornamentals grown for indoor landscaping
Conifers	Ornamentals grown for outdoor landscaping
Deciduous Fruits & Nuts (non-bearing)	

USE SITES - TURF

Athletic Fields	Recreational areas
Golf Courses	Sod Farms
Landscaped areas	Turf around Residential, Institutional, Public,
Lawns	Commercial and Industrial Buildings
Newly seeded turf areas	
Parks	

Active ingredient: *Metalaxyl*
FRAC class: Group 4
Formulation: Liquid
Package size: 2 x 2.5 gallon case

DISEASES CONTROLLED

Downy mildew	Phytophthora spp.
Pythium spp./Yellow tuft	

APPROVED APPLICATION METHODS

Aerial application	Micro sprinkler systems
Drip irrigation systems	Soil drench
Foliar spray	Surface spray on soil
Incorporated in soil mix	Surface spray on turf

USE RATES

Soil surface spray on ornamentals	2 - 5 fl oz/1000 sq ft
Foliar spray on azaleas	1.25 - 2.5 fl oz in 100 gal water
Soil drench	see label*
Soil incorporation	see label*
Established turf	1 - 2 fl oz/1000 sq ft
Newly seeded turf	1 - 2 fl oz/1000 sq ft

* See label for complete directions for use, precautions and restrictions.



Contact your local UPI sales representative or distributor, call 1-800-438-6071 or visit www.upi-usa.com to learn more.

Always read and follow label directions and precautions. Vireo® and UPI logo are registered trademarks of United Phosphorus, Inc. ©November, 2016. UPI, 630 Freedom Business Center, King of Prussia, PA 19406.